arXiv:hep-th/0110029v1 3 Oct 2001

Polarization of the Fulling-Rindler vacuum by uniformly accelerated mirror

Aram A. Saharian*
Department of Physics, Yerevan State University, 1 Alex Manoogian St., 375049 Yerevan, Armenia
(February 1, 2008)

Positive frequency Wightman function and vacuum expectation values of the energy-momentum tensor are computed for a massive scalar field with general curvature coupling parameter and satisfying Robin boundary condition on a uniformly accelerated infinite plate. The both regions of the right Rindler wedge, (i) on the right (RR region) and (ii) on the left (RL region) of the plate are investigated. For the case (ii) the electromagnetic field is considered as well. The mode summation method is used with combination of a variant of the generalized Abel-Plana formula. This allows to present the expectation values in the form of a sum of the purely Rindler and boundary parts. Near the plate surface the vacuum energy-momentum tensor is dominated by the boundary term. At large distances from the plate and near the Rindler horizon the main contribution comes from the purely Rindler part. In the RL region the vacuum energy density of the electromagnetic field is negative near the horizon and is positive near the plate.

PACS number(s): 11.10.Kk, 03.70.+k

I. INTRODUCTION

The influence of boundaries on the vacuum state of a quantum field leads to interesting physical consequences. Well known example is the Casimir effect (see [1–3] and references therein), when the modification of the zero-point fluctuations spectrum by the presence of boundaries induces vacuum forces acting on these boundaries. Another interesting effect is the creation of particles from the vacuum by moving boundaries [4]. In this paper we will study the scalar vacuum polarization brought about by the presence of infinite plane boundary moving by uniform acceleration through the Fulling-Rindler vacuum. This problem for the conformally coupled Dirichlet and Neumann massless scalar and electromagnetic fields in four dimensional spacetime was considered by Candelas and Deutsch [5]. Here we will investigate the vacuum expectation values of the energy-momentum tensor for the massive scalar field with general curvature coupling and satisfying Robin boundary condition on the infinite plane in an arbitrary number of spacetime dimensions. The dimensional dependence of physical quantities is of considerable interest in the Casimir effect and is investigated for various geometries of boundaries and boundary conditions (see [6], [3], [7], [8], [9], [10] and references therein). The Robin boundary conditions are an extension of the ones imposed on perfectly conducting boundaries and may, in some geometries, be useful for depicting the finite penetration of the field into the boundary with the "skin-depth" parameter related to the Robin coefficient [11]. It is interesting to note that the quantum scalar field satisfying Robin condition on the boundary of a cavity violates the Bekenstein's entropy-to-energy bound near certain points in the space of the parameter defining the boundary condition [12]. This type of conditions also appear in considerations of the vacuum effects for a confined charged scalar field in external fields [13] and in quantum gravity [14], [15], [16]. Mixed boundary conditions naturally arise for scalar and fermion bulk fields in the Randall-Sundrum model [17]. In this model the bulk geometry is a slice of anti-de Sitter space and the corresponding Robin coefficient is related to the curvature scale of this space. Unlike to Ref. [5], where the region of the right Rindler wedge on the right from the barrier (in the following we will denote this region as RR) is considered only, we investigate vacuum energy density and stresses in both regions, including the one between the barrier and Rindler horizon (region RL in the following consideration). Our calculation is based on the summation formula derived in Appendix A on the base of the generalized Abel-Plana formula [18] (see also [19]). This allows to extract explicitly the part due to the unbounded Rindler spacetime and to present the boundary part in terms of strongly convergent integrals.

The paper is organized as follows. In the next section we consider the vacuum expectation values for the energy-momentum tensor of the scalar field in the region on the right of the uniformly accelerated infinite plane. By using the summation formula derived in Appendix A these quantities are presented in the form of a sum containing two parts. The first one corresponds to the energy-momentum tensor for the Rindler wedge without boundaries and is investigated in Sec. III. The second term presents the contribution brought about by the presence of the boundary and

---

[*] E-mail: saharyan@server.physdep.r.am

is considered in Sec. IV. The corresponding boundary terms for the region between the barrier and Rindler horizon are evaluated in Sec. V for scalar and electromagnetic fields. In Appendix B we show that the direct evaluation of the energy-momentum tensor in $D = 1$ gives the same result as the analytic continuation from higher dimensions. And finally, in Appendix C, we prove the identity used in Sec. III to see the equivalency of our expressions for the expectation values of the energy-momentum tensor in the case of the Fulling-Rindler vacuum without boundaries to the results previously investigated in the literature for the conformally and minimally coupled scalars.

II. VACUUM EXPECTATION VALUES AND THE WIGHTMAN FUNCTION

Consider a real scalar field $\varphi(x)$ with curvature coupling parameter ζ in D-dimensional space satisfying Robin boundary condition

$$(A + Bn^i\nabla_i)\varphi(x) = 0 \tag{1}$$

on a plane boundary uniformly accelerated normal to itself. Here A and B are constants, n^i is the unit normal to the plane, ∇_i is the covariant derivative operator. The Dirichlet and Neumann boundary conditions are obtained from (1) as special cases. Of course, all results in the following will depend on the ratio A/B only. However, to keep the transition to the Dirichlet and Neumann cases transparent we will use the form (1). The field equation has the form

$$(\nabla_i\nabla^i + m^2 + \zeta R)\varphi(x) = 0, \tag{2}$$

where R is the scalar curvature for the background spacetime. The values of the curvature coupling $\zeta = 0$ and $\zeta = \zeta_c \equiv \frac{D-1}{4D}$ correspond to the minimal and conformal couplings, respectively. In the case of a flat background spacetime the corresponding metric energy-momentum tensor (EMT) is given by (see, e.g., [4])

$$T_{ik} = (1-2\zeta)\partial_i\varphi\partial_k\varphi + (2\zeta - 1/2)g_{ik}\partial^l\varphi\partial_l\varphi - 2\zeta\varphi\nabla_i\nabla_k\varphi + (1/2-2\zeta)m^2g_{ik}\varphi^2. \tag{3}$$

By using the field equation it can be seen that expression (3) can be also presented in the form

$$T_{ik} = \partial_i\varphi\partial_k\varphi + \left[\left(\zeta - \frac{1}{4}\right)g_{ik}\nabla_l\nabla^l - \zeta\nabla_i\nabla_k\right]\varphi^2, \tag{4}$$

and the corresponding trace is equal to

$$T_i^i = D(\zeta - \zeta_c)\nabla_i\nabla^i\varphi^2 + m^2\varphi^2. \tag{5}$$

By virtue of Eq. (4) for the vacuum expectation values (VEV's) of the EMT we have

$$\langle 0\left|T_{ik}(x)\right|0\rangle = \lim_{x'\to x}\nabla_i\nabla'_kG^+(x,x') + \left[\left(\zeta - \frac{1}{4}\right)g_{ik}\nabla_l\nabla^l - \zeta\nabla_i\nabla_k\right]\langle 0\left|\varphi^2(x)\right|0\rangle, \tag{6}$$

where $|0\rangle$ is the amplitude for the vacuum state and we introduced the positive frequency Wightman function:

$$G^+(x,x') = \langle 0\left|\varphi(x)\varphi(x')\right|0\rangle. \tag{7}$$

In Eq. (6) instead of this function one can choose any other bilinear function of fields such as the Hadamard function, Feynman's Green function, etc. The regularized vacuum EMT does not depend on the specific choice. The reason for our choice of the Wightman function is that this function also determines the response of the particle detectors in a given state of motion. By expanding the field operator over eigenfunctions and using the commutation relations one can see that

$$G^+(x,x') = \sum_\alpha \varphi_\alpha(x)\varphi^*_\alpha(x'), \tag{8}$$

where α denotes a set of quantum numbers, $\{\varphi_\alpha(x)\}$ is a complete set of solutions to the field equation (2) satisfying boundary condition (11).

Below we will assume that the plate is situated in the right Rindler wedge, $x_1 > |t|$. It is convenient to introduce Rindler coordinates $(\tau, \xi, \mathbf{x})$ related with Minkowski coordinates $(t, x_1, \mathbf{x})$ by

$$t = \xi\sinh\tau, \quad x_1 = \xi\cosh\tau, \tag{9}$$

where $\mathbf{x} = (x_2, ..., x_D)$ denotes the set of coordinates parallel to the plate. In these coordinates a wordline defined by $\xi, \mathbf{x} = \text{const}$ describes an observer with constant proper acceleration ξ^{-1}. The Minkowski line element restricted to the Rindler wedge is

$$ds^2 = \xi^2 d\tau^2 - d\xi^2 - d\mathbf{x}^{\mathbf{2}}. \tag{10}$$

The corresponding metric is static, admitting the Killing vector field $\partial/\partial\tau$. The Fulling-Rindler vacuum is the vacuum state determined by choosing positive frequency modes to have positive frequency with respect to this Killing vector.

First we will consider the region on the right from the boundary, $\xi \geq a$, RR region, where a^{-1} is the proper acceleration of the barrier. In Rindler coordinates boundary condition (1) for a uniformly accelerated mirror takes the form

$$\left(A + B\frac{\partial}{\partial\xi}\right)\varphi = 0, \quad \xi = a. \tag{11}$$

The problem symmetry allows to write the solutions to field equation (2) as

$$\varphi(x) = C\phi(\xi)e^{i\mathbf{kx} - i\omega\tau}. \tag{12}$$

Substituting this into Eq. (2) we see that the function $\phi(\xi)$ satisfies the equation

$$\xi\frac{d}{d\xi}\left(\xi\frac{d\phi}{d\xi}\right) + \left(\omega^2 - \xi^2\lambda^2\right)\phi(\xi) = 0, \quad \lambda = \sqrt{k^2 + m^2}. \tag{13}$$

The corresponding linearly independent solutions are the Bessel modified functions $I_{\pm i\omega}(\lambda\xi)$ and $K_{i\omega}(\lambda\xi)$ with the imaginary order. It can be seen that for any two solutions to equation (13), $\phi_\omega^{(1)}(\xi)$ and $\phi_\omega^{(2)}(\xi)$, the following integration formula takes place

$$\int \frac{d\xi}{\xi}\phi_\omega^{(1)}(\xi)\phi_{\omega'}^{(1)}(\xi) = \frac{\xi}{\omega^2 - \omega'^2}\left[\phi_\omega^{(1)}(\xi)\frac{d\phi_{\omega'}^{(2)}(\xi)}{d\xi} - \phi_{\omega'}^{(2)}(\xi)\frac{d\phi_\omega^{(1)}(\xi)}{d\xi}\right]. \tag{14}$$

As we will see below the normalization integrals for the eigenfunctions will have this form.

For the region $\xi > a$ a complete set of solutions that are of positive frequency with respect to $\partial/\partial\tau$ and bounded as $\xi \to \infty$ is

$$\varphi_\alpha(x) = CK_{i\omega}(\lambda\xi)e^{i\mathbf{kx} - i\omega\tau}, \quad \alpha = (\omega, \mathbf{k}). \tag{15}$$

From boundary condition (11) we find that the possible values for ω have to be roots to the equation

$$AK_{i\omega}(\lambda a) + B\lambda K'_{i\omega}(\lambda a) = 0, \tag{16}$$

where the prime denotes the differentiation with respect to the argument. This equation has infinite number of real zeros. We will denote them by $\omega = \omega_n = \omega_n(k)$, $n = 1, 2, ...$ arranged in ascending order: $\omega_n < \omega_{n+1}$. The coefficient C in (15) is determined by the normalization condition

$$(\varphi_\alpha, \varphi_{\alpha'}) = -i\int d\mathbf{x}\int\frac{d\xi}{\xi}\varphi_\alpha \overset{\leftrightarrow}{\partial}_\tau \varphi_{\alpha'}^* = \delta_{\alpha\alpha'} \tag{17}$$

with respect to the standard Klein-Gordon inner product, and the ξ-integration goes over the region (a, ∞). Using integration formula (14) with $\phi_\omega^{(1)} = \phi_\omega^{(2)} = K_{i\omega}(\lambda\xi)$ and Wronskian $W\{K_\nu(z), I_\nu(z)\} = 1/z$, from (17) one finds

$$C^2 = \frac{\pi}{(2\pi)^D\omega_n}\left[\int_a^\infty \frac{d\xi}{\xi}K_{i\omega_n}^2(\lambda\xi)\right]^{-1} = \frac{1}{(2\pi)^{D-1}}\frac{\bar{I}_{i\omega_n}(\lambda a)}{\frac{\partial}{\partial\omega}\bar{K}_{i\omega}(\lambda a)\mid_{\omega=\omega_n}}, \tag{18}$$

where for a given function $f(z)$ we use the notation

$$\bar{f}(z) = Af(z) + bzf'(z) = 0, \quad b = B/a. \tag{19}$$

From the first equality in Eq. (18) it follows that C^2 is real and positive. Now substituting the eigenfunctions (15) into (8) for the Wightman function we obtain

$$G^{+}(x,x') = \int \frac{d\mathbf{k}}{(2\pi)^{D-1}}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \sum_{n=1}^{\infty} \frac{\bar{I}_{i\omega}(\lambda a)}{\frac{\partial}{\partial\omega}\bar{K}_{i\omega}(\lambda a)} K_{i\omega}(\lambda\xi) K_{i\omega}(\lambda\xi') e^{-i\omega(\tau-\tau')}|_{\omega=\omega_n}. \quad (20)$$

For the further evolution of VEV's (20) we can apply to the sum over n summation formula (A5). As a function $F(z)$ in this formula let us choose

$$F(z) = K_{iz}(\lambda\xi) K_{iz}(\lambda\xi') e^{-iz(\tau-\tau')}. \quad (21)$$

Using the asymptotic formulas for the Bessel modified functions it can be easily seen that condition (A3) is satisfied if $a^2 e^{|\tau-\tau'|} < |\xi\xi'|$. Note that this condition is satisfied in the coincidence limit $\tau = \tau'$ for the points in the region under consideration, $\xi, \xi' > a$. With $F(z)$ from (21) the contribution corresponding to the integral term on the left of formula (A5) is the Wightman function for the Fulling-Rindler vacuum $|0_R\rangle$ without boundaries:

$$G^{+}_R(x,x') = \langle 0_R|\varphi(x)\varphi(x')|0_R\rangle = \frac{1}{\pi^2} \int \frac{d\mathbf{k}}{(2\pi)^{D-1}}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^{\infty} d\omega \sinh(\pi\omega) e^{-i\omega(\tau-\tau')} K_{i\omega}(\lambda\xi) K_{i\omega}(\lambda\xi'). \quad (22)$$

To see this note that for the right Rindler wedge the eigenfunctions to the field equation (2) are in form (see, for instance, [20], [21], [22], [23])

$$\varphi_{\omega\mathbf{k}} = \frac{\sqrt{\sinh \pi\omega}}{(2\pi)^{(D-1)/2}\pi} e^{i\mathbf{k}\mathbf{x} - i\omega\tau} K_{i\omega}(\lambda\xi). \quad (23)$$

Substituting these modes into formula (8) we obtain Eq. (22). Hence, taking into account this and applying summation formula (A5) to Eq. (20) we receive

$$G^{+}(x,x') = G^{+}_R(x,x') + \langle\varphi(x)\varphi(x')\rangle^{(b)}, \quad (24)$$

where the second term on the right is induced by the existence of the barrier:

$$\langle\varphi(x)\varphi(x')\rangle^{(b)} = -\frac{1}{\pi} \int \frac{d\mathbf{k}}{(2\pi)^{D-1}}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^{\infty} d\omega \frac{\bar{I}_{\omega}(\lambda a)}{\bar{K}_{\omega}(\lambda a)} K_{\omega}(\lambda\xi) K_{\omega}(\lambda\xi') \cosh[\omega(\tau-\tau')], \quad (25)$$

and is finite for $\xi > a$. All divergences in the coincidence limit are contained in the first term corresponding to the Fulling - Rindler vacuum without boundaries. The VEV's for the EMT are obtained substituting Eq. (24) into Eq. (6). Similar to (24) this VEV's are also in the form of the sum containing purely Rindler and boundary parts. Firstly we will concentrate to the first term.

III. VACUUM EMT IN THE RINDLER WEDGE WITHOUT BOUNDARIES

The Wightman function for the right Rindler wedge given by formula (22) is divergent in the coincidence limit $x' \to x$. This leads to the divergences in the VEV for the EMT. To extract these divergences we can subtract from Eq. (22) the corresponding Wightman function for the Minkowski vacuum $|0_M\rangle$. By using mode sum formula (8) and the standard Minkowskian eigenfunctions it easily can be seen that the latter may be presented in the form

$$G^{+}_M(x,x') = \int \frac{d^{D-1}\mathbf{k}}{(2\pi)^{D}} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} K_0\left(\lambda\sqrt{\xi^2 + \xi'^2 - 2\xi\xi' \cosh(\tau-\tau')}\right), \quad (26)$$

where $\xi^2 + \xi'^2 - 2\xi\xi' \cosh(\tau-\tau') = (x_1 - x_1')^2 - (t-t')^2$. Using the integration formula [24]

$$K_0\left(\lambda\sqrt{\xi^2 + \xi'^2 + 2\xi\xi' \cos b}\right) = \frac{2}{\pi} \int_0^{\infty} d\omega\, \cosh(b\omega) K_{i\omega}(\lambda\xi) K_{i\omega}(\lambda\xi'), \quad (27)$$

Eq. (26) can be presented in the form convenient for subtraction:

$$G^{+}_M(x,x') = \frac{1}{\pi^2} \int \frac{d^{D-1}\mathbf{k}}{(2\pi)^{D-1}} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^{\infty} d\omega\, \cosh\left\{\omega\left[\pi - i(\tau-\tau')\right]\right\} K_{i\omega}(\lambda\xi) K_{i\omega}(\lambda\xi'). \quad (28)$$

As a result for the difference between the Rindler and Minkowskian Wightman functions one obtains

$$G_R^+(x,x') - G_M^+(x,x') = -\frac{1}{\pi^2}\int \frac{d^{D-1}\mathbf{k}}{(2\pi)^{D-1}} e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \times \int_0^\infty d\omega\, e^{-\pi\omega} \cos\left[\omega(\tau-\tau')\right] K_{i\omega}(\lambda\xi) K_{i\omega}(\lambda\xi'). \tag{29}$$

After integrating over directions of the vector $\mathbf{k}$, in particular, for the difference between VEV's of the field square it follows from here

$$\begin{aligned}\langle\varphi^2(x)\rangle_{\rm sub}^{(R)} &= \langle 0_R|\varphi(x)|0_R\rangle - \langle 0_M|\varphi(x)|0_M\rangle \\ &= \frac{-1}{2^{D-2}\pi^{(D+3)/2}\Gamma\left(\frac{D-1}{2}\right)}\int_0^\infty dk\, k^{D-2}\int_0^\infty d\omega\, e^{-\pi\omega} K_{i\omega}^2(\lambda\xi).\end{aligned} \tag{30}$$

Using formula (6), for the corresponding difference between the VEV's of the EMT we receive

$$\begin{aligned}\langle T_i^k\rangle_{\rm sub}^{(R)} &= \langle 0_R|T_i^k|0_R\rangle - \langle 0_M|T_i^k|0_M\rangle \\ &= \frac{-\delta_i^k}{2^{D-2}\pi^{(D+3)/2}\Gamma\left(\frac{D-1}{2}\right)}\int_0^\infty dk k^{D-2}\lambda^2\int_0^\infty d\omega e^{-\pi\omega} f^{(i)}\left[K_{i\omega}(\lambda\xi)\right],\end{aligned} \tag{31}$$

where we have introduced the following notations

$$\begin{aligned}f^{(0)}[g(z)] &= \left(\frac{1}{2}-2\zeta\right)\left|\frac{dg(z)}{dz}\right|^2 + \frac{\zeta}{z}\frac{d}{dz}\left|g(z)\right|^2 + \left[\frac{1}{2}-2\zeta+\left(\frac{1}{2}+2\zeta\right)\frac{\omega^2}{z^2}\right]\left|g(z)\right|^2, \\ f^{(1)}[g(z)] &= -\frac{1}{2}\left|\frac{dg(z)}{dz}\right|^2 - \frac{\zeta}{z}\frac{d}{dz}\left|g(z)\right|^2 + \frac{1}{2}\left(1-\frac{\omega^2}{z^2}\right)\left|g(z)\right|^2, \\ f^{(i)}[g(z)] &= \left(\frac{1}{2}-2\zeta\right)\left[\left|\frac{dg(z)}{dz}\right|^2 + \left(1-\frac{\omega^2}{z^2}\right)\left|g(z)\right|^2\right] - \frac{k^2}{(D-1)\lambda^2}\left|g(z)\right|^2, \quad i=2,3,...,D,\end{aligned} \tag{32}$$

and indices 0 and 1 correspond to the coordinates τ and ξ.

We consider first the massless limit. In this limit interchanging the orders of integration, making use of the integration formula [24]

$$\mathcal{I}_D(\xi) \equiv \int_0^\infty dk\, k^D K_{i\omega}^2(k\xi) = \frac{2^{D-2}}{\xi^{D+1}\Gamma(D+1)}\Gamma^2\left(\frac{D+1}{2}\right)\left|\Gamma\left(\frac{D+1}{2}+i\omega\right)\right|^2 \tag{33}$$

and the relations

$$\int_0^\infty dk\, k^{D-1} K_{i\omega}(k\xi) K'_{i\omega}(k\xi) = -\frac{D-1}{2\xi}\mathcal{I}_{D-2}(\xi), \tag{34}$$

$$\int_0^\infty dk\, k^D K'^2_{i\omega}(k\xi) = \frac{\mathcal{I}_{D-2}(\xi)}{4D\xi^2}\left[4\omega^2+(D+1)(D-1)^2\right], \tag{35}$$

for the purely Rindler parts (30) and (31) one finds

$$\langle\varphi^2\rangle_{\rm sub}^{(R)} = -\frac{\xi^{-D+1}}{2^D\pi^{D/2+1}\Gamma(D/2)}\int_0^\infty d\omega\, e^{-\pi\omega}\left|\Gamma\left(\frac{D-1}{2}+i\omega\right)\right|^2, \tag{36}$$

$$\langle T_i^k\rangle_{\rm sub}^{(R)} = -\frac{\delta_i^k\xi^{-D-1}}{2^D\pi^{D/2+1}\Gamma(D/2)}\int_0^\infty d\omega\, \omega^2 e^{-\pi\omega}\left|\Gamma\left(\frac{D-1}{2}+i\omega\right)\right|^2 f_0^{(i)}(\omega), \tag{37}$$

where we have introduced notations

$$\begin{aligned}f_0^{(0)}(\omega) &= -Df_0^{(1)}(\omega) = 1 + D(D-1)(\zeta_c-\zeta)/\omega^2, \\ f_0^{(i)}(\omega) &= -1/D + (D-1)^2(\zeta_c-\zeta)/\omega^2, \quad i=2,3,\ldots,D.\end{aligned} \tag{38}$$

Using the recurrence formula, the gamma function in Eqs. (36) and (37) can be led to $\Gamma(1+i\omega)$ for odd D and to $\Gamma(1/2+i\omega)$ for even D. The standard formulas for the latters [25] allow us to write expressions (36) and (37) in the form

$$\langle\varphi^2\rangle_{\rm sub}^{(R)} = -\frac{\xi^{-D+1}}{2^{D-1}\pi^{D/2}\Gamma(D/2)}\int_0^\infty \frac{\omega^{D-2}d\omega}{e^{2\pi\omega}+(-1)^D}\prod_{l=1}^{l_m}\left[\left(\frac{D-1-2l}{2\omega}\right)^2+1\right], \tag{39}$$

$$\langle T_i^k\rangle_{\rm sub}^{(R)} = -\frac{\delta_i^k\xi^{-D-1}}{2^{D-1}\pi^{D/2}\Gamma(D/2)}\int_0^\infty \frac{\omega^{D}d\omega}{e^{2\pi\omega}+(-1)^D}f_0^{(i)}(\omega)\prod_{l=1}^{l_m}\left[\left(\frac{D-1-2l}{2\omega}\right)^2+1\right], \tag{40}$$

where $l_m = D/2-1$ for even $D>2$ and $l_m=(D-1)/2$ for odd $D>1$, and the value for the product over l is equal to 1 for $D=1,2,3$. These relations illustrate the thermal properties of the Minkowski vacuum relative to the Rindler observer [20], [21], [22]. As we see from Eq. (40), though thermal with temperature $T=(2\pi\xi)^{-1}$, the spectrum of the vacuum EMT, in general, has a non-Planckian form: the density of states factor is not proportional to $\omega^{D-1}d\omega$. It is interesting to note that for even values of the space dimension D this thermal distribution for the scalar field is Fermi-Dirac type.

Note that for the case of a two dimensional spacetime we can not directly put $D=1$ into the integral (40). Now, though the factor $D-1$ in the second term on the right of expression (38) for $f_0^{(0)}(\omega)$ is equal zero, the corresponding ω-integral diverges and this term gives finite contribution to the energy density. Keeping $D>1$ the ω-integral in Eq. (40) can be expressed in terms of the Riemann zeta function $\zeta_R(x)$, and in the limit $D\to 1$ one obtains

$$\langle T_0^0\rangle_{\rm sub}^{(R)} = -\langle T_1^1\rangle_{\rm sub}^{(R)} = -\frac{1}{\pi\xi^2}\left(\frac{1}{24}+\lim_{D\to 1}\Gamma(D)\zeta_R(D-1)\right) = \frac{1}{2\pi\xi^2}\left(\zeta-\frac{1}{12}\right), \tag{41}$$

where $\zeta_R(0)=-1/2$.

The density of states factor is Planckian for the conformally coupled scalar and for $D=1,2,3$:

$$\langle T_i^k\rangle_{\rm sub}^{(R)} = -\frac{\xi^{-D-1}}{2^{D-1}\pi^{D/2}\Gamma(D/2)}\int_0^\infty \frac{\omega^{D}d\omega}{e^{2\pi\omega}+(-1)^D}{\rm diag}\left(1,-\frac{1}{D},...,-\frac{1}{D}\right), \quad \zeta=\zeta_c, \quad D=1,2,3. \tag{42}$$

This result for $D=3$ case was obtained by Candelas and Deutsch [5]. Expression (42) corresponds to the absence from the vacuum of thermal distribution. This means that the Minkowski vacuum corresponds to a thermal state with respect to uniformly accelerated observers [22].

We now treat the case of a massive scalar field. First of all let us present the difference (29) between Rindler and Minkowski Wightman functions in another alternative form. For this we will follow the procedure already used in Ref. [23] for the Green function. Substituting [26]

$$K_{i\omega}(\lambda\xi)K_{i\omega}(\lambda\xi') = \frac{1}{2}\int_{-\infty}^{+\infty}dy\, e^{i\omega y}K_0(\lambda\gamma), \tag{43}$$

where $\gamma^2=\xi^2+\xi'^2+2\xi\xi'\cosh y$, into Eq. (29) and integrating over ω we receive

$$G_R^+(x,x')-G_M^+(x,x') = -\int\frac{d^{D-1}\mathbf{k}}{(2\pi)^D}e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')}\int_{-\infty}^{+\infty}dy\,\frac{K_0(\lambda\gamma)}{\pi^2+(y+\tau-\tau')^2}. \tag{44}$$

Integrating over directions of the vector $\mathbf{k}$ with the help of formula

$$\int d^{D-1}\mathbf{k}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')}F(k) = (2\pi)^{(D-1)/2}\int_0^\infty dk\, k^{D-2}F(k)\frac{J_{(D-3)/2}(k\,|\mathbf{x}-\mathbf{x}'|)}{(k\,|\mathbf{x}-\mathbf{x}'|)^{(D-3)/2}}, \tag{45}$$

with $J_\nu(x)$ being the Bessel function, and using the value for the integral over k [24],

$$\int_0^\infty dk\, k^{\mu+1}J_\mu(bk)K_0(\gamma\sqrt{k^2+m^2}) = b^\mu m^{\mu+1}K_{\mu+1}(m\sqrt{b^2+m^2}), \tag{46}$$

one obtains

$$G_R^+(x,x')-G_M^+(x,x') = -\frac{1}{2\pi}\int_{-\infty}^{+\infty}\frac{dy}{\pi^2+y^2}\left(\frac{m}{2\pi\gamma_1}\right)^{(D-1)/2}K_{(D-1)/2}(m\gamma_1), \tag{47}$$

where $\gamma_1^2=\xi^2+\xi'^2+2\xi\xi'\cosh(y-\tau+\tau')+|\mathbf{x}-\mathbf{x}'|^2$. The similar relation for the Feynman's Green function is considered in Ref. [23]. Noting that γ_1 is just the geodesic separation of the points $x=(\tau,\xi,\mathbf{x})$ and $x''=(\tau'+y,-\xi',\mathbf{x})$

and taking into account the standard expression for the Minkowskian Wightman function, relation (47) can be written as

$$G_R^+(x,x') = G_M^+(x,x') - \int_{-\infty}^{+\infty} \frac{dy}{\pi^2+y^2} G_M^+(x,x''). \tag{48}$$

Following Ref. [23], the second term on the right of this formula we can interpret as coming from an image charge density $-1/(\pi^2+y^2)$ distributed on a line $(-\xi',\mathbf{x})$ in the left Rindler wedge.

In the coincidence limit Eq. (47) yields

$$\langle \varphi^2(x)\rangle_{\rm sub}^{(R)} = -\frac{2m^{D-1}}{(2\pi)^{(D+1)/2}} \int_0^\infty \frac{dy}{\pi^2+y^2} \frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}}, \quad z = 2m\xi\cosh(y/2). \tag{49}$$

This quantity is monotone increasing negative function on ξ. Substituting Eqs. (47) and (49) into formula (6) for the difference between the corresponding VEV's of the EMT we obtain (no summation over i)

$$\langle T_0^0\rangle_{\rm sub}^{(R)} = -\frac{2m^{D+1}}{(2\pi)^{(D+1)/2}} \int_0^\infty \frac{dy}{y^2+\pi^2} \left[\frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}} + D\frac{K_{(D+1)/2}(z)}{z^{(D+1)/2}}\right] \left(1-4\zeta\cosh^2(y/2)\right), \tag{50}$$

$$\langle T_1^1\rangle_{\rm sub}^{(R)} = \frac{2m^{D+1}}{(2\pi)^{(D+1)/2}} \int_0^\infty \frac{dy}{y^2+\pi^2} \frac{K_{(D+1)/2}(z)}{z^{(D+1)/2}} \left(1-4\zeta\cosh^2(y/2)\right), \tag{51}$$

$$\begin{aligned} \langle T_i^i\rangle_{\rm sub}^{(R)} = {} & \frac{2m^{D+1}}{(2\pi)^{(D+1)/2}} \int_0^\infty \frac{dy}{y^2+\pi^2} \left\{\frac{K_{(D+1)/2}(z)}{z^{(D+1)/2}} + \right. \\ & \left. + \left[\frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}} + (D-1)\frac{K_{(D+1)/2}(z)}{z^{(D+1)/2}}\right](4\zeta-1)\cosh^2(y/2)\right\}, \quad i=2,3,\ldots,D. \end{aligned} \tag{52}$$

Note that near the horizon, in the limit $m\xi \ll 1$, the leading terms of the asymptotic expansion for the vacuum EMT components (50)-(52) do not depend on the mass and coincide with Eq. (40). Hence, near the horizon these components diverge as ξ^{-D-1}. The VEV's of the EMT for the real massive scalar fields with minimal and conformal couplings on background of the Rindler spacetime were considered in Ref. [27] by using the covariant functional Schrödinger formalism, developed in Ref. [28]. In this paper for the EMT the form (3) is used with $\zeta=0$ and $\zeta=\zeta_c$ and the corresponding formulas [formulas (2.39) and (2.45) in Ref. [27]] are more complicated. The equivalency of these formulas to our expressions (50)-(52) with the special values $\zeta=0,\zeta_c$ can be seen using the identity

$$\begin{aligned} \frac{4}{m^2\xi^2}\int_0^\infty dy \frac{\pi^2-3y^2}{(\pi^2+y^2)^3} \frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}} = {} & \int_0^\infty \frac{dy}{\pi^2+y^2}\left\{(1-\cosh y)\frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}}\right. \\ & \left. + \left[D+1-(D-1)\cosh y\right]\frac{K_{(D+1)/2}(z)}{z^{(D+1)/2}}\right\}, \end{aligned} \tag{53}$$

where z is defined as in Eq. (49). The proof of this identity we give in Appendix C.

From expressions (49)-(52) it follows that for the massive scalar field the energy spectrum is not strictly thermal and the quantities $\langle 0_M\left|\varphi^2\right|0_M\rangle - \langle 0_R\left|\varphi^2\right|0_R\rangle = -\langle\varphi^2(x)\rangle_{\rm sub}^{(R)}$ and $\langle 0_M\left|T_i^k\right|0_M\rangle - \langle 0_R\left|T_i^k\right|0_R\rangle = -\langle T_i^k(x)\rangle_{\rm sub}^{(R)}$ do not coincide with the corresponding ones for the Minkowski thermal bath, $\langle\varphi^2\rangle_M^{\rm th}$ and $\langle T_i^k\rangle_M^{\rm th}$, with temperature $(2\pi\xi)^{-1}$ (for this statement in a variety of situations see Refs. [29]- [32] and references therein). To illustrate this for the case $D=3$ we present in Fig. 1 the quantities $-\langle\varphi^2\rangle_{\rm sub}^{(R)}$ (curve a) and

$$\langle\varphi^2\rangle_M^{\rm th} = \frac{1}{2\pi^2}\int_0^\infty \frac{\left(\omega^2+2m\omega\right)^{1/2}d\omega}{e^{(m+\omega)/T}-1}, \quad T = \frac{1}{2\pi\xi} \tag{54}$$

(curve b) in dependence of $m\xi$.



FIG. 1. The $D=3$ expectation values for the field square in the case of the Rindler vacuum without boundaries, $-\xi^2\langle\varphi^2\rangle_{\rm sub}^{(R)}$ (curve a) and for the Minkowski thermal bath, $\xi^2\langle\varphi^2\rangle_M^{\rm th}$ (curve b), given by formula (54), versus $m\xi$.

The integrals in formulas (50)-(52) are strongly convergent and useful for numerical evaluation. In Fig. 2 the results of the corresponding numerical evaluation for the vacuum EMT components are presented in the Rindler case without boundaries for minimally and conformally coupled scalars.



FIG. 2. The expectation values for the diagonal components of the energy-momentum tensor, $\xi^{D+1}\langle T_i^i\rangle^{(R)}$ (no summation over i), versus $m\xi$ in the case of the Fulling-Rindler vacuum without boundaries for minimally (left) and conformally (right) coupled scalar fields in $D=3$. The curves a, b, c correspond to the values $i=0,1,2$ respectively.

In the limit $m\xi\gg 1$ using the asymptotic formula for the Macdonald function and taking into account that the main contribution into the integrals in Eqs. (50), (51), and (52) comes from the region with small y, in the leading order one finds

$$\langle T_0^0\rangle_{\rm sub}^{(R)}\sim\langle T_i^i\rangle_{\rm sub}^{(R)}\sim -2m\xi\langle T_1^1\rangle_{\rm sub}^{(R)}\sim\frac{(4\zeta-1)m^{(D+1)/2}}{2^D\pi^{(D+3)/2}\xi^{(D+1)/2}}e^{-2m\xi},\quad i=2,...,D,\quad m\xi\to\infty. \tag{55}$$

In the massless limit, $m=0$, from Eq. (49) for the field product one obtains

$$\langle\varphi^2(x)\rangle_{\rm sub}^{(R)}=-\frac{\Gamma\left(\frac{D-1}{2}\right)}{2^D\pi^{(D+1)/2}\xi^{D-1}}\int_0^\infty\frac{dy}{\pi^2+y^2}\cosh^{1-D}(y/2). \tag{56}$$

The corresponding formulas for the VEV's of the EMT are written as

$$\langle T_0^0\rangle_{\rm sub}^{(R)}=-D\langle T_1^1\rangle_{\rm sub}^{(R)}=\frac{D\Gamma\left(\frac{D+1}{2}\right)}{(2\sqrt{\pi}\xi)^{D+1}}\int_0^\infty\frac{dy}{y^2+\pi^2}\frac{4\zeta\cosh^2(y/2)-1}{\cosh^{D+1}(y/2)}, \tag{57}$$

$$\langle T_i^i\rangle_{\rm sub}^{(R)}=\frac{\Gamma\left(\frac{D+1}{2}\right)}{(2\sqrt{\pi}\xi)^{D+1}}\int_0^\infty\frac{dy}{y^2+\pi^2}\frac{1-(1-4\zeta)(D-1)\cosh^2(y/2)}{\cosh^{D+1}(y/2)},\quad i=2,\ldots,D. \tag{58}$$

Note that the relation (57) between the energy density and the vacuum effective pressure also follows from the continuity equation for the EMT (see Eq. (63) below). Formulas (56)-(58) are alternative representations for the VEV's (36), (37).

IV. VEV'S IN THE RR REGION

In Sec. II we have seen that the Wightman function in the case of the presence of a barrier can be presented in the form (24), where the first summand on the right corresponds to the Fulling-Rindler vacuum without boundaries and the second one is induced by the barrier and is given by expression (25). In the coincidence limit from this formula for the boundary part of the field square we have

$$\langle\varphi^2(x)\rangle^{(b)} = \frac{-1}{2^{D-2}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)}\int_0^\infty dk\, k^{D-2}\int_0^\infty d\omega\, \frac{\bar{I}_\omega(\lambda a)}{\bar{K}_\omega(\lambda a)}K_\omega^2(\lambda\xi), \quad \xi > a. \tag{59}$$

This quantity is monotone increasing negative function on ξ for Dirichlet scalar and monotone decreasing positive function for Neumann scalar. Substituting the Wightmann function (24) into Eq. (6) and taking into account Eqs. (25) and (59) for the VEV's of the EMT in the region $\xi > a$ one finds

$$\langle 0|T_i^k(x)|0\rangle = \langle 0_R|T_i^k(x)|0_R\rangle + \langle T_i^k(x)\rangle^{(b)}, \tag{60}$$

where the first term on the right is the VEV for the Fulling-Rindler vacuum without boundaries and is investigated in previous section. The second term is the contribution brought by the presence of the barrier at $\xi = a$:

$$\langle T_i^k\rangle^{(b)} = \frac{-\delta_i^k}{2^{D-2}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)}\int_0^\infty dk k^{D-2}\lambda^2\int_0^\infty d\omega \frac{\bar{I}_\omega(\lambda a)}{\bar{K}_\omega(\lambda a)}F^{(i)}\left[K_\omega(\lambda\xi)\right], \tag{61}$$

where the functions $F^{(i)}[g(z)]$, $i = 0, 1, 2$ are obtained from the functions $f^{(i)}[g(z)]$ in Eqs. (32) replacing $\omega \to i\omega$:

$$F^{(i)}[g(z)] = f^{(i)}[g(z), \omega \to i\omega]. \tag{62}$$

It can be easily checked that both summands on the right of Eq. (60) satisfy the continuity equation $T_{i;k}^k = 0$, which for the geometry under consideration takes the form

$$\frac{d}{d\xi}(\xi T_1^1) - T_0^0 = 0. \tag{63}$$

By using the inequalities

$$I_\omega'(z) < \sqrt{1+\omega^2/z^2}I_\omega(z), \quad -K_\omega'(z) > \sqrt{1+\omega^2/z^2}K_\omega(z) \tag{64}$$

it can be easily seen that $F^{(0)}[K_\omega(z)] > 0$, $F^{(1)}[K_\omega(z)] < 0$ for $\zeta \leq 0$ and $F^{(i)}[K_\omega(z)] > 0$, $i = 2, 3, \ldots, D$ for $\zeta \leq \frac{D-3}{4(D-1)}$. In the cases of Dirichlet and Neumann scalars for the boundary parts of the EMT components these leads

$$\left(\langle T_0^0\rangle^{(b)}\right)_{\text{Dirichlet}} < 0, \quad \left(\langle T_1^1\rangle^{(b)}\right)_{\text{Dirichlet}} > 0 \tag{65}$$

$$\left(\langle T_0^0\rangle^{(b)}\right)_{\text{Neumann}} < 0, \quad \left(\langle T_1^1\rangle^{(b)}\right)_{\text{Neumann}} > 0 \tag{66}$$

for $\zeta \leq 0$, and (no summation over i)

$$\left(\langle T_i^i\rangle^{(b)}\right)_{\text{Dirichlet}} < 0, \quad \left(\langle T_i^i\rangle^{(b)}\right)_{\text{Neumann}} > 0, \quad i = 2, 3, \ldots, D \tag{67}$$

for $\zeta \leq \frac{D-3}{4(D-1)}$.

For $D = 1$ case the corresponding boundary part is obtained from Eq. (61)

$$\langle T_i^k\rangle^{(b)} = -\frac{\delta_i^k}{\pi}m^2\int_0^\infty d\omega \frac{\bar{I}_\omega(ma)}{\bar{K}_\omega(ma)}F^{(i)}\left[K_\omega(m\xi)\right], \quad i = 0, 1. \tag{68}$$

Taking the limit $m \to 0$ for the massless field we receive from here

$$\langle T_i^k \rangle^{(b)} = \frac{\zeta \delta_i^k}{\pi \xi^2} \int_0^\infty d\omega \, \frac{A + B\omega/a}{A - B\omega/a} e^{-2\omega \ln(\xi/a)} F_0^{(i)}, \tag{69}$$

with

$$F_0^{(0)} = 2\omega + 1, \quad F_0^{(1)} = -1. \tag{70}$$

In this case our assumption for the absence of imaginary eigenmodes corresponds to the restriction $AB \leq 0$ on the Robin coefficients. We have obtained result (69) by using the analytical continuation from the values $D > 1$. If we want to realize a direct evaluation in $D = 1$ the prescription described in Sec. II has to be modified for the massless case. In appendix B we show that the corresponding evaluation leads to the same result.

Boundary part (61) is finite for all values $\xi > a$ and diverges at the plate surface $\xi = a$. These surface divergences are well known in quantum field theory with boundaries and are investigated for various type of boundary conditions and geometries. To extract the leading part of the boundary divergence note that near the boundary the main contribution into the ω-integral comes from large values of ω and we can use the uniform asymptotic expansions for the modified Bessel functions [25]. Introducing a new integration variable $k \to \omega k$ and replacing the Bessel modified functions by their uniform asymptotic expansions in the leading order one obtains

$$\langle T_0^0(x) \rangle^{(b)} \sim \langle T_2^2(x) \rangle^{(b)} \sim \frac{D(2\delta_{B0} - 1)(\zeta - \zeta_c)}{2^D \pi^{(D+1)/2} (\xi - a)^{D+1}} \Gamma\left(\frac{D+1}{2}\right), \quad \xi \to a \tag{71}$$

$$\langle T_1^1(x) \rangle^{(b)} \sim \frac{(1 - 2\delta_{B0})(\zeta - \zeta_c)}{2^D \pi^{(D+1)/2} a (\xi - a)^D} \Gamma\left(\frac{D+1}{2}\right). \tag{72}$$

These terms do not depend on mass and Robin coefficients, and have opposite signs for the Dirichlet and Neumann boundary conditions. Note that the leading terms for $\langle T_0^0(x) \rangle^{(b)}$ and $\langle T_2^2(x) \rangle^{(b)}$ are the same as for the plate in Minkowski spacetime (see, for instance, [8]). This statement is also valid for the spherical and cylindrical boundaries on background of the Minkowski spacetime [9], [10].

Now let us consider the asymptotic behavior of the boundary part (61) for large ξ, $\xi \gg a$. Introducing in Eq. (61) a new integration variable $y = \lambda \xi$ and using the asymptotic formulas for the Bessel modified functions for small values of the argument, we see that the subintegrand is proportional to $(ya/2\xi)^{2\omega}$. It follows from here that the main contribution into the ω-integral comes from the small values of ω. Expanding with respect to ω in the leading order we obtain

$$\langle T_i^k(x) \rangle^{(b)} \sim -\frac{\delta_i^k (-1)^{\delta_{A0}} \xi^{-D-1} A_0^{(i)}(m\xi)}{2^D 3 \pi^{(D-1)/2} (1 + \delta_{A0}) \Gamma\left(\frac{D-1}{2}\right) \ln^2(2\xi/a)}, \tag{73}$$

where

$$A_0^{(i)}(m\xi) = \int_{m\xi}^\infty dy \, y^3 (y^2 - m^2\xi^2)^{(D-3)/2} F^{(i)}[K_\omega(y)]|_{\omega=0}. \tag{74}$$

If, in addition, one has $m\xi \gg 1$ the integral in this formula can be evaluated replacing the Macdonald function by its asymptotic for large values of the argument. In the leading order this yields

$$A_0^{(0)} \sim A_0^{(i)} \sim -2m\xi A_0^{(1)} \sim \pi(1/4 - \zeta) \Gamma\left(\frac{D-1}{2}\right) (m\xi)^{(D+1)/2} e^{-2m\xi}, \quad m\xi \gg 1, \quad i = 2, 3, \cdots, D. \tag{75}$$

For the massless case the integral in Eq. (74) may be evaluated using formulas (33)–(35) and one obtains

$$A_0^{(0)} \sim -DA_1^{(1)} \sim \frac{D}{D-1} A_2^{(2)} \sim \frac{2^D D(\zeta_c - \zeta)}{(D-1)^2 \Gamma(D)} \Gamma^4\left(\frac{D+1}{2}\right), \quad m = 0. \tag{76}$$

For the conformally coupled scalar the leading terms vanish and the VEV's are proportional to $\xi^{-D-1} \ln^{-3}(2\xi/a)$ for $\xi/a \gg 1$. As it follows from Eqs. (40), (55), and (73) in both cases (75) and (76) and in the region far from the boundary the VEV's (60) are dominated by the purely Fulling-Rindler part (the first summand on the right): $\langle T_i^k \rangle^{(b)} / \langle T_i^k \rangle_{\text{sub}}^{(R)} \sim \ln^{-2}(2\xi/a)$. From Eq. (73) we see that for a given ξ the boundary part tends to zero as $a \to 0$ (the barrier coincides with the right Rindler horizon) and the corresponding VEV's of the EMT are the same as for the Fulling-Rindler vacuum without boundaries. Hence, the barrier located at the Rindler horizon does not alter the vacuum EMT.

And finally, we turn to the asymptotic $a, \xi \to \infty$, $\xi - a = \mathrm{const}$. In this limit $\xi/a \to 1$ and ω-integrals in Eq. (61) are dominated by the large ω. Replacing the modified Bessel functions by their uniform asymptotic expansions, keeping the leading terms only and introducing a new integration variable $\nu = \omega/a$ for the energy density one obtains

$$\begin{aligned}\langle T_0^0\rangle^{(b)} \sim & -\frac{1}{2^{D-1}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)}\int_0^\infty dk\, k^{D-2}\int_0^\infty d\nu \frac{e^{-2(\xi-a)\sqrt{\nu^2+\lambda^2}}}{\sqrt{\nu^2+\lambda^2}} \\ & \times \frac{A+B\sqrt{\nu^2+\lambda^2}}{A-B\sqrt{\nu^2+\lambda^2}}\left[-4\zeta\nu^2+\lambda^2(1-4\zeta)\right]. \end{aligned} \tag{77}$$

Converting to polar coordinates, $\nu = r\cos\theta$, $k = r\sin\theta$, we can integrate over angles with the help of the formula

$$\int_0^{\pi/2} d\theta\, \sin^n\theta = \frac{\sqrt{\pi}}{2}\frac{\Gamma\left(\frac{n+1}{2}\right)}{\Gamma(n/2+1)}. \tag{78}$$

Introducing the new integration variable $y = \sqrt{r^2+m^2}$ we receive

$$\begin{aligned}\langle 0|T_0^0|0\rangle \sim \langle T_0^0\rangle^{(b)} \sim & \frac{1}{2^D\pi^{D/2}\Gamma(D/2)}\int_m^\infty dy\, (y^2-m^2)^{D/2}e^{-2(\xi-a)y} \\ & \times\frac{A+By}{A-By}\left[4(\zeta-\zeta_c)+\frac{m^2(4\zeta-1)}{y^2-m^2}\right], \quad a,\xi\to\infty, \quad \xi-a=\mathrm{const}. \end{aligned} \tag{79}$$

Here we have taken into account that in the limit $\xi \to \infty$ the purely Rindler part $\langle 0_R|T_0^0|0_R\rangle$ vanishes and in Eq. (60) the boundary term remains only. In the same order the evaluation of the components, corresponding to the vacuum effective pressures in the plane parallel to the barrier, leads (no summation over i)

$$\langle 0|T_i^i|0\rangle \sim \langle 0|T_0^0|0\rangle, \quad i = 2, 3, ..., D, \tag{80}$$

The leading term for the component $\langle 0|T_1^1|0\rangle$ vanishes,

$$\langle 0|T_1^1|0\rangle = \mathcal{O}(1/a), \quad a \to \infty. \tag{81}$$

The term to the next order can be found using continuity equation (63) and expression (79). The expectation values (79), (80), and (81) coincide with the corresponding VEV's induced by a single plate in the Minkowski spacetime with the Robin boundary conditions on it, and are investigated in Ref. [8] for the massless field and in Ref. [33] for the massive scalar field.

V. VACUUM STRESSES IN THE RL REGION

A. Scalar field

In previous section we have considered VEV's for the EMT in the region $\xi > a$. Now we will turn to the case of the scalar field in region between the barrier and Rindler horizon (RL region), $0 < \xi < a$, and satisfying boundary condition (11) on the boundary $\xi = a$. For this region we have to take both solutions to equation (13) and the eigenfunctions have the form (12), where now

$$\phi(\xi) = Z_{i\omega}(\lambda\xi, \lambda a) \equiv K_{i\omega}(\lambda\xi) - \frac{\bar{K}_{i\omega}(\lambda a)}{\bar{I}_{i\omega}(\lambda a)}I_{i\omega}(\lambda\xi), \tag{82}$$

and the coefficient is obtained from boundary condition (11). Here the quantities with overbars are defined in accordance with Eq. (19). As we see, unlike to the previous case, the spectrum for ω is continous. The normalization coefficient is determined from condition (17), where integration over ξ goes between the limits $(0, a)$. Using integration formula (14) and the representation for the delta function

$$\delta(x) = \lim_{\sigma\to\infty}\frac{\sin\sigma x}{\pi x}, \tag{83}$$

it can be seen that

$$C = \frac{\sqrt{\sinh \omega\pi}}{\pi (2\pi)^{(D-1)/2}}. \tag{84}$$

Substituting eigenfunctions (12) with the function (82) into the mode sum for the Wightman function one finds

$$G^{+}(x,x') = \frac{1}{\pi^2} \int \frac{d^{D-1}\mathbf{k}}{(2\pi)^{D-1}}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^\infty d\omega \sinh(\pi\omega) e^{-i\omega(\tau-\tau')} Z_{i\omega}(\lambda\xi, \lambda a) Z^{*}_{i\omega}(\lambda\xi', \lambda a). \tag{85}$$

The part induced by the barrier can be obtained subtracting from here the Wightman function for the right Rindler wedge without boundaries, given by expression (22). To transform the corresponding difference note that

$$Z_{i\omega}(\lambda\xi, \lambda a) Z^{*}_{i\omega}(\lambda\xi', \lambda a) - K_{i\omega}(\lambda\xi) K^{*}_{i\omega}(\lambda\xi') = \frac{\pi \bar{K}_{i\omega}(\lambda a)}{2i \sinh \pi\omega} \sum_{\sigma=-1,1} \sigma \frac{I_{i\sigma\omega}(\lambda\xi) I_{i\sigma\omega}(\lambda\xi')}{\bar{I}_{i\sigma\omega}(\lambda a)}, \tag{86}$$

where we have used the standard relation

$$K_\nu(z) = \frac{\pi}{2\sin \nu\pi} \left[I_{-\nu}(z) - I_\nu(z) \right]. \tag{87}$$

This allows to present the boundary part in the form

$$\begin{aligned} \langle \varphi(x)\varphi(x')\rangle^{(b)} &= G^{+}(x,x') - G^{+}_{R}(x,x') \\ &= \frac{1}{i} \int \frac{d^{D-1}\mathbf{k}}{(2\pi)^{D}}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^\infty d\omega e^{-i\omega(\tau-\tau')} \sum_{\sigma=-1,1} \sigma \frac{\bar{K}_{i\omega}(\lambda a)}{\bar{I}_{i\sigma\omega}(\lambda a)} I_{i\sigma\omega}(\lambda\xi) I_{i\sigma\omega}(\lambda\xi'). \end{aligned} \tag{88}$$

Assuming that the function $\bar{I}_{i\omega}(\lambda a)$ ($\bar{I}_{-i\omega}(\lambda a)$) has no zeros for $-\pi/2 \leq \arg\omega < 0$ ($0 < \arg\omega < \pi/2$) we can rotate the integration contour over ω by angle $-\pi/2$ for the term with $\sigma = 1$ and by angle $\pi/2$ for the term with $\sigma = -1$. The integrals taken around the arcs of large radius tend to zero under the condition $|\xi\xi'| < a^2 e^{|\tau-\tau'|}$ (note that, in particular, this is the case in the coincidence limit for the region under consideration). As a result for the difference of the Wightman functions one obtains

$$\langle \varphi(x)\varphi(x')\rangle^{(b)} = -\frac{1}{\pi} \int \frac{d^{D-1}\mathbf{k}}{(2\pi)^{D-1}}\, e^{i\mathbf{k}(\mathbf{x}-\mathbf{x}')} \int_0^\infty d\omega \frac{\bar{K}_\omega(\lambda a)}{\bar{I}_\omega(\lambda a)} I_\omega(\lambda\xi) I_\omega(\lambda\xi') \cosh\left[\omega(\tau-\tau')\right]. \tag{89}$$

As we see this expression differs from the corresponding formula (25) for the region $a < \xi < \infty$ by replacements $I_\omega \to K_\omega$, $K_\omega \to I_\omega$. In the coincidence limit for the VEV of the field square we receive

$$\langle \varphi^2(x)\rangle^{(b)} = \frac{-1}{2^{D-2}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)} \int_0^\infty dk\, k^{D-2} \int_0^\infty d\omega \frac{\bar{K}_\omega(\lambda a)}{\bar{I}_\omega(\lambda a)} I^2_\omega(\lambda\xi), \quad 0 < \xi < a. \tag{90}$$

This quantity is monotone decreasing negative function on ξ for Dirichlet scalar and monotone increasing positive function for Neumann scalar. From Eq. (6) the VEV's of the EMT are obtained in the form (60) with the boundary part

$$\langle T_i^k\rangle^{(b)} = -\frac{\delta_i^k}{2^{D-2}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)} \int_0^\infty dk\, k^{D-2} \lambda^2 \int_0^\infty d\omega\, \frac{\bar{K}_\omega(\lambda a)}{\bar{I}_\omega(\lambda a)} F^{(i)}\left[I_\omega(\lambda\xi)\right], \tag{91}$$

where, as in Eq. (61), the functions $F^{(i)}[g(z)]$ are obtained from expressions (32) replacing $\omega \to i\omega$ (see Eq. (62)). Comparing Eqs. (90), (91), (59), and (61) we see that the boundary parts for the region $0 < \xi < a$ can be obtained from the corresponding ones for the region $\xi > a$ by the replacements $I_\omega \to K_\omega$, $K_\omega \to I_\omega$. Note that here the situation is the same as for the interior and exterior regions in the case of cylindrical and spherical surfaces on background of the Minkowski spacetime (see, [9], [10]). The reason for this analogy is that the uniformly accelerated observers produce world lines in Minkowski spacetime which in Euclidean space would correspond to circles. This correspondence extends also to the field equation and its solutions, and modes (12) are the Minkowski spacetime analogue corresponding to cylinder harmonics.

For $D = 1$ case the integral over $\mathbf{k}$ in Eq. (89) is absent and $\lambda = m$. The corresponding formulas for the field square and vacuum EMT are obtained from Eqs. (90) and (91) replacing

$$\frac{1}{2^{D-2}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)}\int_0^\infty dk\, k^{D-2} \to \frac{1}{\pi}, \quad \lambda \to m. \tag{92}$$

In particular, for the massless $D=1$ case we have

$$\langle T_i^k\rangle^{(b)} = \frac{\zeta\delta_i^k}{\pi\xi^2}\int_0^\infty d\omega\, \frac{A - B\omega/a}{A + B\omega/a} e^{-2\omega\ln(a/\xi)}\tilde{F}_0^{(i)}, \tag{93}$$

where

$$\tilde{F}_0^{(0)} = 2\omega - 1, \quad \tilde{F}_0^{(1)} = 1. \tag{94}$$

The expressions (91) are divergent on the plate surface, $\xi = a$. The leading terms for the corresponding asymptotic expansions are derived by the way similar to that for Eqs. (71) and (72), and can be obtained from these formulas replacing $\xi - a$ by $a - \xi$ and changing the sign for $\langle T_1^1\rangle^{(b)}$.

Now let us consider the behavior of the boundary part (91) for $\xi \ll a$. Introducing instead of k a new integration variable $y = \lambda a$ and using the formula for $I_\omega(y\xi/a)$ in the case of small values of the argument we see that the ω -subintegrand is proportional to $e^{-2\omega\ln(a/y\xi)}$. It follows from here that the main contribution comes from the small values of ω. Expanding over ω we obtain the standard integrals of the form $\int_0^\infty \omega^n e^{-2\omega\ln(a/y\xi)}d\omega$. Performing these integrals one obtains in the leading order

$$\langle T_0^0\rangle^{(b)} \sim -\langle T_1^1\rangle^{(b)} \sim -\frac{\zeta B_0(ma)\left(1 + \mathcal{O}(\ln^{-1}(2a/\xi))\right)}{2^{D-1}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)a^{D-1}\xi^2\ln^2(2a/\xi)} \tag{95}$$

$$\langle T_i^i\rangle^{(b)} \sim \frac{(4\zeta - 1)B_0(ma)\left(1 + \mathcal{O}(\ln^{-1}(2a/\xi))\right)}{2^{D}\pi^{(D+1)/2}\Gamma\left(\frac{D-1}{2}\right)a^{D-1}\xi^2\ln^3(2a/\xi)}, \quad i = 2, 3, ..., D, \tag{96}$$

where

$$B_0(ma) = \int_{ma}^\infty dy\, y(y^2 - m^2a^2)^{(D-3)/2}\frac{\bar{K}_0(y)}{\bar{I}_0(y)}. \tag{97}$$

Note that the function $B_0(ma)$ is positive for Dirichlet boundary condition and is negative for Neumann one. For $D=1$ case one has

$$\langle T_0^0\rangle^{(b)} \sim -\langle T_1^1\rangle^{(b)} \sim -\frac{\zeta}{2\pi\xi^2\ln^2(2/m\xi)}\frac{\bar{K}_0(ma)}{\bar{I}_0(ma)}. \tag{98}$$

As we see the boundary part is divergent at Rindler horizon. Recall that near the horizon the purely Rindler part behaves as ξ^{-D-1} and, therefore, in this limit the total vacuum EMT is dominated by this part.

B. Electromagnetic field

We now turn to the case of the electromagnetic field in the region $0 < \xi < a$ for the case $D = 3$. We will assume that the mirror is a perfect conductor with the standard boundary conditions of vanishing of the normal component of the magnetic field and the tangential components of the electric field, evaluated at the local inertial frame in which the conductor is instantaneously at rest. As it has been shown in Ref. [5] the corresponding eigenfunctions for the vector potential $A_\mu(x)$ may be resolved into transverse electric (TE) and transverse magnetic (TM) (with respect to ξ-direction) modes:

$$A_\mu^{(\sigma)}(x) = \left\{ \begin{array}{ll} (0, 0, -ik_3, ik_2)\varphi^{(0)}(x), & \text{for} \quad \sigma = 0 \\ (-\xi\partial/\partial\xi, i\omega/\xi, 0, 0)\varphi^{(1)}(x), & \text{for} \quad \sigma = 1 \end{array} \right. , \tag{99}$$

where $\mathbf{k} = (k_2, k_3)$, $\sigma = 0$ and $\sigma = 1$ correspond to the TE and TM waves respectively. From the perfect conductor boundary conditions on the vector potential we obtain the corresponding boundary conditions for the scalar fields $\varphi^{(\sigma)}(x)$:

$$\varphi^{(0)}(x) = 0, \quad \frac{\partial\varphi^{(1)}(x)}{\partial\xi} = 0, \quad \xi = a. \tag{100}$$

As a result the TE/TM modes correspond to the Dirichlet/Neumann scalars. In the corresponding expressions for the eigenfunctions $A_{\alpha\mu}^{(\sigma)}(x)$ the normalization coefficient is determined from the orthonormality condition

$$\int d\mathbf{x} \int_0^a \frac{d\xi}{\xi} A_\alpha^{(\sigma)\mu}(x) A_{\alpha'\mu}^{(\sigma')*}(x) = -\frac{2\pi}{\omega}\delta_{\alpha\alpha'}\delta_{\sigma\sigma'}, \quad \alpha = (\mathbf{k}, \omega). \tag{101}$$

From here for the corresponding Diriclet and Neumann modes we have

$$\varphi_\alpha^{(\sigma)}(x) = \frac{\sqrt{\sinh\omega\pi}}{\pi^{3/2}k}\left[K_{i\omega}(k\xi) - \frac{K_{i\omega}^{(\sigma)}(ka)}{I_{i\omega}^{(\sigma)}(ka)} I_{i\omega}(k\xi)\right] e^{i\mathbf{kx}-i\omega\tau}, \tag{102}$$

where $K_{i\omega}^{(0)}(z) = K_{i\omega}(z)$, $K_{i\omega}^{(1)}(z) = dK_{i\omega}(z)/dz$ and the same notations for the function $I_{i\omega}(z)$. The VEV's for the EMT can be obtained substituting eigenfunctions (99) into the mode sum formula

$$\langle 0 \left| T_i^k(x) \right| 0\rangle = \sum_{\sigma=0,1} \int d\mathbf{k} \int_0^\infty d\omega\, T_i^k \left\{ A_{\alpha\mu}^{(\sigma)}(x), A_{\alpha\mu}^{(\sigma)*}(x) \right\}, \tag{103}$$

with the standard bilinear form for the electromagnetic field EMT:

$$T_i^k \left\{ A_\mu(x), A_\mu(x) \right\} = \frac{1}{4\pi}\left(-F_{il}F^{lk} + \frac{1}{4}\delta_i^k F_{ln}F^{ln} \right), \tag{104}$$

where $F_{lk} = \partial A_k/\partial x^l - \partial A_l/\partial x^k$ is the field tensor. Substituting eigenfunctions (99) into the mode sum (103) one finds

$$\langle 0 \left| T_i^k(x) \right| 0\rangle = \frac{\delta_i^k}{4\pi^3} \sum_{\sigma=0,1} \int_0^\infty dk\, k^3 \int_0^\infty d\omega\, \sinh(\omega\pi) f_{\rm em}^{(i)} \left[K_{i\omega}(k\xi) - \frac{K_{i\omega}^{(\sigma)}(ka)}{I_{i\omega}^{(\sigma)}(ka)} I_{i\omega}(k\xi) \right]. \tag{105}$$

Here for a given function $g(z)$ the following notations are introduced

$$\begin{aligned} f_{\rm em}^{(0)}[g(z)] &= \left| \frac{dg(z)}{dz} \right|^2 + \left(1 + \frac{\omega^2}{z^2} \right) |g(z)|^2, \\ f_{\rm em}^{(1)}[g(z)] &= -\left| \frac{dg(z)}{dz} \right|^2 + \left(1 - \frac{\omega^2}{z^2} \right) |g(z)|^2, \\ f_{\rm em}^{(2)}[g(z)] &= f_{\rm em}^{(3)}[g(z)] = -|g(z)|^2. \end{aligned} \tag{106}$$

It can be easily checked that components (105) satisfy zero trace condition and continuity equation (63). The way for subtraction from these quantities the parts due to the Fulling-Rindler vacuum without boundaries is the same as for the scalar field, given above in this section. This yields to the following result

$$\langle 0 \left| T_i^k \right| 0\rangle = \langle 0_R \left| T_i^k \right| 0_R\rangle - \frac{\delta_i^k}{4\pi^2} \int_0^\infty dk\, k^3 \int_0^\infty d\omega \left[\frac{K_\omega(ka)}{I_\omega(ka)} + \frac{K'_\omega(ka)}{I'_\omega(ka)} \right] F_{\rm em}^{(i)}[I_\omega(k\xi)], \tag{107}$$

where

$$\langle 0_R \left| T_i^k \right| 0_R\rangle = \langle 0_M \left| T_i^k \right| 0_M\rangle - \frac{1}{\pi^2\xi^4} \int_0^\infty \frac{d\omega\,(\omega^3+\omega)}{e^{2\pi\omega}-1} {\rm diag}\left(1, -\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3} \right) \tag{108}$$

are the VEV's for the Fulling-Rindler vacuum without boundaries [5], and the functions $F_{\rm em}^{(i)}[g(z)]$ are obtained from Eq. (106) replacing $\omega \to i\omega$: $F_{\rm em}^{(i)}[g(z)] = f_{\rm em}^{(i)}[g(z), \omega \to i\omega]$. Note that the ω-integral in Eq. (108) is equal to 11/240. Comparing the boundary part in Eq. (107) to the corresponding expression for the RR region, $\xi > a$, derived in [5], we see that, as in the scalar case, these results are obtained from each other by the replacements $I_\omega \to K_\omega$, $K_\omega \to I_\omega$.

By using inequalities (64) we conclude that

$$\frac{K_\omega(z)}{I_\omega(z)} + \frac{K'_\omega(z)}{I'_\omega(z)} < 0. \tag{109}$$

This, together with the observation that $I_\omega(z)$ is monotone increasing function, immediately shows that the components $\langle T_i^i \rangle^{(b)}$, $i = 2, 3$ are negative and monotone decreasing on ξ. Further from the first inequality (64) one has $F_{\rm em}^{(1)}[I_\omega(k\xi)] > 0$ and it can be seen that this function is monotone increasing on ξ. Hence, the component $\langle T_1^1 \rangle^{(b)}$ is positive and monotone increasing. And, at last, by using Eq. (64) we see that $F_{\rm em}^{(0)}[I_\omega(k\xi)] > 0$, and as a result one obtains that the boundary part of the energy density is positive, $\langle T_0^0 \rangle^{(b)} > 0$.

Now let us consider the asymptotics of the boundary part in Eq. (107) near the barrier, $\xi \to a$. In this limit the corresponding expressions are divergent. It follows from here that the main contributions into the VEV's are due to the large ω. Rescaling the integration variable $k \to \omega k$ and replacing the Bessel modified functions by their uniform asymptotic expansions in the leading order one finds

$$\begin{aligned} \langle T_0^0 \rangle^{(b)} &\sim -2\langle T_i^i \rangle^{(b)} \sim \frac{1}{30\pi^2 a(a-\xi)^3}, \quad i = 2, 3, \\ \langle T_1^1 \rangle^{(b)} &\sim \frac{1}{60\pi^2 a^2 (a-\xi)^2}, \quad \xi \to a - 0. \end{aligned} \tag{110}$$

Note that the corresponding asymptotics for the region $\xi > a$ are given by the same formulas [5].

Now we turn to the asymptotics in the limit $\xi \to 0$. Replacing the function $I_\omega(k\xi)$ by its asymptotic, and noting that the main contribution into the ω-integrals come from the region with small ω, in the leading order we have (no summation over i)

$$\langle T_0^0 \rangle^{(b)} \sim \langle T_1^1 \rangle^{(b)} \sim -\langle T_i^i \rangle^{(b)} \sim -\frac{1}{8\pi^2 a^4 \ln(2a/\xi)} \int_0^\infty dy\, y^3 \left[\frac{K_0(y)}{I_0(y)} - \frac{K_1(y)}{I_1(y)} \right]. \tag{111}$$

In this formula the value for the integral is equal to -1.326. As we see, unlike to the case for the scalar field, the boundary part of the electromagnetic vacuum EMT tends to zero as $\xi \to 0$.

And finally, let us consider the leading terms in the vacuum EMT in the limit $a, \xi \to \infty$, $\xi - a =$ constant. Introducing in Eq. (107) a new integration variable ka we see that in this limit the corresponding terms have the same form as in the case $\xi \to a$, $a =$ constant and are given by expressions (110). Note that in this limit the total vacuum EMT is dominated by the boundary part.

In Fig. 3 we have plotted the electromagnetic vacuum energy density, $a^4 \langle T_0^0 \rangle$ (left), and perpendicular pressure, $-a^4 \langle T_1^1 \rangle$ (right), in dependence of ξ/a, generated by a single perfectly conducting plate uniformly accelerated normal to itself. For the plate $\xi = a$. The dashed curves present the corresponding regularized quantities for the Fulling-Rindler vacuum without boundaries, given by the second summand on the right of formula (108). The full curves are for the boundary parts, second term on the right of formula (107). In addition to the region $0 < \xi < a$, investigated in this section, we have also included the region $a < \xi < \infty$. The formulas for the boundary part in the latter case are derived in Ref. [5] and, as it have been mentioned above, are obtained from the ones considered here by replacements $I_\omega \to K_\omega$, $K_\omega \to I_\omega$. From Fig. 3 we see that the vacuum perpendicular pressure is always negative.



FIG. 3. The expectation values for the boundary part of the energy density, $a^4 \langle T_0^0 \rangle^{(b)}$ (left), and perpendicular effective pressure, $-a^4 \langle T_1^1 \rangle^{(b)}$ (right), multiplied by a^4, for the electromagnetic field versus ξ/a. The dashed curves present the corresponding regularized quantities for the Fulling-Rindler vacuum without boundaries [see Eq. (108)].

The total energy density (as a sum of purely Rindler and boundary parts) is always negative in the region $a < \xi < \infty$. For the region $0 < \xi < \infty$ the total energy density is negative near the horizon and is positive near the plate. As a result for some intermediate value of ξ the energy density vanishes.

VI. CONCLUSIONS

Quantum field theory in accelerated systems contains many of the special features produced by a gravitational field without the complications due to a curved background spacetime. In this paper we have considered the VEV's of the EMT induced by uniformly accelerated plane boundary in the Fulling-Rindler vacuum. This problem for the region on the right of the plane has been previously studied by Candelas and Deutch in the paper [5]. These authors consider a conformally coupled scalar field with Dirichlet or Neumann boundary conditions and electromagnetic field. Here we generalize the corresponding results for (i) an arbitrary number of spatial dimension, (ii) for a scalar field with general curvature coupling parameter and (iii) nonzero mass, (iv) for mixed boundary conditions of Robin type. In addition, (v) we consider the VEV's in the region between the plane and Rindler horizon for scalar and electromagnetic fields.

To obtain the expectation values for the energy-momentum tensor we first construct the positive frequency Wightman function (this function is also important in considerations of the response of a particle detector [4]). For the region on the right of the plate, $\xi > a$, the application of the generalized Abel-Plana formula to the mode sum over zeros of the function $\bar{K}_{i\omega}$ allows us to extract the purely Rindler part without boundaries and to present the additional boundary part in terms of strongly convergent integrals (formulas (24), (25)). The expectation values for the EMT are obtained by applying a certain second order differential operator and taking the coincidence limit. First, in Sec. III, we consider the vacuum EMT in the Rindler wedge without boundaries. After the subtraction of the Minkowskian part the corresponding components are presented in the form (31) with notations (32). For the massless scalar the VEV's of the EMT take the form (40). This expression corresponds to the absence from the vacuum of thermal distribution with standard temperature $T = (2\pi\xi)^{-1}$. In general, this distribution has non-Planckian spectrum: the density of states factor is not proportional to $\omega^{D-1}d\omega$. The spectrum takes the Planckian form for conformally coupled scalars in $D = 1, 2, 3$. It is interesting to note that for even values of spatial dimension the distribution is Fermi-Dirac type. Further, on the base of procedure already used in [23], for the massive case we present the vacuum EMT in another alternative form, formulas (50)-(52). By using the identity proved in Appendix C we show that for the cases of conformally and minimally coupled scalars these formulas are equivalent to those previously obtained in Ref. [27]. For the massive scalar the energy spectrum is not strictly thermal and the corresponding quantities do not coincide with ones for the Minkowski thermal bath with temperature $(2\pi\xi)^{-1}$. This fact is illustrated in Fig. 1. In the limit $m\xi \gg 1$ the vacuum EMT components are exponentially decreasing functions on $m\xi$. In the massless limit we obtain another equivalent representations, Eqs. (57) and (58).

In Sec. IV we investigate the boundary part of the vacuum EMT induced by a single plate in the region $\xi > a$, formula (61). The corresponding result for the massless 1D field can be obtained by the analytic continuation and has form (69). In Appendix B we show that this result coincides with the formula derived by direct calculation in $D = 1$. Further we investigate the asymptotic properties of the boundary EMT. Near the plate surface the total VEV's are dominated by the boundary part and the corresponding components diverge at the boundary. For non-conformally coupled scalars the leading terms are given by formulas (71) and (72) and are the same as for an infinite plane boundary in Minkowski spacetime with Dirichlet and Neumann boundary conditions. These terms do not depend on the mass or Robin coefficients, and have opposite signs for Dirichlet and Neumann cases. For large values of $\xi \gg a$ the leading term in the asymptotic expansion of the boundary induced EMT does not depend on the Robin coefficient B and has the form (73). In this limit $\langle T_i^k\rangle^{(b)}/\langle T_i^k\rangle_{\rm sub}^{(R)} \sim \ln^{-2}(2\xi/a)$ and the vacuum EMT is dominated by the purely Rindler part. For a given ξ the boundary part of the EMT tends to zero as $a \to 0$, i.e. $\langle T_i^k\rangle^{(b)} \to 0$ in the limit when the barrier coincides with the Rindler horizon. As a result the corresponding VEV's of the EMT are the same as for the Fulling-Rindler vacuum without boundaries. Hence, the barrier located at the Rindler horizon does not alter the vacuum EMT.

The vacuum stresses in the region $0 < \xi < a$ are investigated in Sec. V for scalar and electromagnetic fields. The corresponding boundary parts can be presented as Eqs. (91) and (107), respectively. These formulas differ from the ones for the region $\xi > a$ by the replacements $I_\omega \to K_\omega, K_\omega \to I_\omega$. For the scalar field the boundary part diverges at the horizon with the leading behavior (95), (96). The divergence of the purely Rindler part is stronger and this part dominates near the Rindler horizon. Unlike to the scalar case, the boundary parts of the vacuum EMT components for the electromagnetic field vanish at the horizon (see asymptotic formulas (111)). In Fig. 3 we present the vacuum densities for the electromagnetic field, generated by a perfectly conducting plate in both RR and RL regions. As seen, unlike to the RR region, where the energy density is negative for all $a < \xi < \infty$, in the RL region the energy density is negative near the Rindler horizon, but is positive near the plate. As a result for some intermediate value of ξ we have zero energy density.

ACKNOWLEDGEMENTS

I am grateful to Professor E. Chubaryan and Professor A. Mkrtchyan for general encouredgement and suggestions, and to R. Avagyan, L. Grigoryan and A. Yeranyan for useful discussions. This work was supported by the Armenian National Science and Education Fund (ANSEF) Grant No. PS14-00.

APPENDIX A: SUMMATION FORMULA OVER ZEROS OF $\bar{K}_\omega$

In Sec. II we have shown that the VEV's for the EMT components in the RR region contain sums over zeros $\omega = \omega_n(\eta)$, $n = 1, 2, \ldots$ of the function $\bar{K}_{i\omega}(\eta) = AK_{i\omega}(\eta) + bzK'_{i\omega}(\eta)$ for a given η. To obtain a summation formula over these zeros we will use the generalized Abel-Plana formula derived in [18], [19]. In this formula as functions $f(z)$ and $g(z)$ let us choose

$$\begin{aligned} f(z) &= \frac{2i}{\pi}\sinh \pi z\, F(z), \\ g(z) &= \frac{\bar{I}_{iz}(\eta) + \bar{I}_{-iz}(\eta)}{\bar{K}_{iz}(\eta)} F(z), \end{aligned} \tag{A1}$$

with a function $F(z)$ analytic in the right half-plane $\mathrm{Re}\, z \geq 0$, and quantities with overbars are defined in accordance with Eq. (19). For the sum and difference of these functions one has

$$g(z) \pm f(z) = 2F(z)\frac{\bar{I}_{\mp iz}(\eta)}{\bar{K}_{iz}(\eta)}. \tag{A2}$$

By using the asymptotic formulas for the Bessel modified functions for large values of index, the conditions for the generalized Abel-Plana formula can be written in terms of the function $F(z)$ as follows

$$|F(z)| < \epsilon(|z|)e^{-\pi x}\left(\frac{|z|}{\eta}\right)^{2|y|}, \quad z = x + iy, \quad x > 0, \quad |z| \to \infty, \tag{A3}$$

where $|z|\epsilon(|z|) \to 0$ when $|z| \to \infty$. Let $\omega = \omega_n$ be zeros for the function $\bar{K}_{i\omega}(\eta)$ in the right half-plane. Here we will assume values of A and b for which all these zeros are real. It can be seen that they are simple. This directly follows from the relation

$$\frac{\partial}{\partial \omega}\bar{K}_{i\omega}(\eta)\bigg|_{\omega=\omega_n} = \frac{2\omega_n b}{K_{i\omega_n}(\eta)}\int_1^\infty \frac{d\xi}{\xi}K^2_{i\omega_n}(\xi\eta)\ , \tag{A4}$$

which is a direct consequence of integral formula (14) with $\phi^{(1)}_\omega(\xi) = \phi^{(2)}_\omega(\xi) = K_{i\omega_n}(\xi\eta)$. Substituting functions (A1) into generalized Abel-Plana formula (formula (2.11) in Ref. [19]) and taking into account that the points $z = \omega_n$ are simple poles for the function $g(z)$ and the relation $\bar{I}_{i\omega_n}(\eta) = \bar{I}_{-i\omega_n}(\eta)$, one obtains

$$\begin{aligned} &\lim_{l\to\infty}\left\{\sum_{n=1}^{n_l}\frac{\bar{I}_{i\omega_n}(\eta)F(\omega_n)}{\partial\bar{K}_{i\omega}(\eta)/\partial\omega|_{\omega=\omega_n}} - \frac{1}{\pi^2}\int_0^l \sinh \pi z\, F(z)dz\right\} \\ &\qquad = -\frac{1}{2\pi}\int_0^\infty \frac{\bar{I}_z(\eta)}{\bar{K}_z(\eta)}\left[F(ze^{\pi i/2}) + F(ze^{-\pi i/2})\right]dz, \end{aligned} \tag{A5}$$

where $\omega_{n_l} < l < \omega_{n_l+1}$. Here we have assumed that the function $F(z)$ is analytic in the right half-plane. However, this formula can be easily generalized for the functions having poles in this region.

APPENDIX B: $D = 1$ MASSLESS CASE: DIRECT EVALUATION

Here we show that the direct evaluation of the boundary VEV's in $D = 1$ gives the same result as the analytical continuation from the higher values of the space dimension. Now in equation (13) for the function $\phi(\xi)$ one has $\lambda = 0$. The corresponding linearly independent solutions are $e^{\pm i\omega \ln(\xi/a)}$. The normalized eigenfunctions satisfying boundary condition (11) are in form

$$\varphi_\omega(\xi, \tau) = \frac{e^{-i\omega\tau}}{\sqrt{\pi\omega}} \sin\left[\omega \ln(\xi/a) - \beta\right], \quad 0 < \omega < \infty, \tag{B1}$$

where

$$e^{i\beta} = \frac{A + i\omega B/a}{\sqrt{A^2 + (\omega B/a)^2}}. \tag{B2}$$

Note that these eigenfunctions have the same form for the regions $\xi > a$ and $\xi < a$. Unlike to $D > 1$ cases now the both types of the eigenfunctions are bounded and the spectrum for ω in the region $\xi > a$ is continous. Substituting eigenfunctions (B1) into the mode sum formula (8) after some algebra one finds the Wightman function in the form (24), where the first summand on the right is the $D = 1$ Wightman function for the Rindler wedge without boundaries:

$$G_R^+(x, x') = \frac{1}{2\pi} \int_0^\infty \frac{d\omega}{\omega} e^{-i\omega(\tau-\tau')} \cos\left[\omega \ln(\xi/\xi')\right], \tag{B3}$$

and for the boundary part one has

$$\langle\varphi(x)\varphi(x')\rangle^{(b)} = -\frac{1}{2\pi} \int_0^\infty \frac{d\omega}{\omega} e^{-i\omega(\tau-\tau')} \cos\left[\omega \ln(\xi\xi'/a^2) - 2\beta\right]. \tag{B4}$$

To transform this expression we write the cos function in terms of exponentials and rotate the integration contour by angle $\pm\pi/2$ for the term with $\exp[i\omega \ln(\xi\xi'/a^2)]$ and by angle $\mp\pi/2$ for the term with $\exp[-i\omega \ln(\xi\xi'/a^2)]$. Here and below the upper and lower signs correspond to the cases $\xi, \xi' > a$ and $\xi, \xi' < a$ respectively. Under the condition $|\tau - \tau'| < |\ln(\xi\xi'/a^2)|$ the integrals over the arcs with large radius in the complex ω-plane tend to zero and we receive

$$\langle\varphi(x)\varphi(x')\rangle^{(b)} = -\frac{1}{2\pi} \int_0^\infty \frac{d\omega}{\omega} \frac{A \pm \omega B/a}{A \mp \omega B/a} e^{-\omega|\ln(\xi\xi'/a^2)|} \cosh[\omega(\tau - \tau')]. \tag{B5}$$

The boundary contribution to the vacuum EMT can be found substituting this expression into formula (6). It can be easily seen that as a result for the boundary part we obtain formulas (69) and (93). Hence, we have shown that the direct evaluation gives the same result as the analytic continuation.

APPENDIX C: PROOF OF IDENTITY (53)

In this appendix we will prove identity (53), which was used to see the equivalency of our formulas (50)-(52) for the VEV's of the EMT in the special cases of the conformally and minimally coupled massive scalars to the results derived by Hill in Ref. [27]. Our starting point is the integral

$$\mathcal{L}(\xi, \xi') = \int_0^\infty dk\, k^{D-2} \int_0^\infty d\omega\, \omega^2 e^{-\pi\omega} K_{i\omega}(\lambda\xi) K_{i\omega}(\lambda\xi') \tag{C1}$$

with λ defined as in Eq. (13). By using formulas (43) and [24]

$$\int_0^\infty dk\, k^{D-2} K_0\left(\gamma\sqrt{k^2 + m^2}\right) = 2^{(D-3)/2} m^{D-1} \Gamma\left(\frac{D-1}{2}\right) \frac{K_{(D-1)/2}(m\gamma)}{(m\gamma)^{(D-1)/2}}, \tag{C2}$$

for this integral one obtains

$$\mathcal{L}(\xi, \xi') = 2^{(D-1)/2} \pi m^{D-1} \Gamma\left(\frac{D-1}{2}\right) \int_0^\infty dy\, \frac{\pi^2 - 3y^2}{(\pi^2 + y^2)^3} \frac{K_{(D-1)/2}(m\gamma)}{(m\gamma)^{(D-1)/2}}, \tag{C3}$$

where γ is defined by Eq. (43). In particular taking $\xi' = \xi$ we receive

$$\mathcal{L}(\xi,\xi) = 2^{(D-1)/2}\pi m^{D-1}\Gamma\left(\frac{D-1}{2}\right)\int_0^\infty dy\, \frac{\pi^2-3y^2}{(\pi^2+y^2)^3}\frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}}, \tag{C4}$$

with z defined in accordance to Eq.(49). Another form for $\mathcal{L}(\xi,\xi')$ can be obtained by using the relation

$$\omega^2 K_{i\omega}(\lambda\xi) = \lambda^2\xi^2 K_{i\omega}(\lambda\xi) - \xi^2\frac{d^2K_{i\omega}(\lambda\xi)}{d\xi^2} - \xi\frac{dK_{i\omega}(\lambda\xi)}{d\xi}. \tag{C5}$$

The substitution into Eq. (C1) yields

$$\begin{aligned}\mathcal{L}(\xi,\xi') = 2^{(D-3)/2}\pi m^{D+1}\Gamma\left(\frac{D-1}{2}\right)\int_0^\infty \frac{dy}{\pi^2+y^2}\left\{(D-1)\frac{K_{(D+1)/2}(m\gamma)}{(m\gamma)^{(D+1)/2}}\right. \\ \left.+\left[1-\frac{d^2}{d(m\xi)^2}-\frac{1}{m\xi}\frac{d}{d(m\xi)}\right]\frac{K_{(D-1)/2}(m\gamma)}{(m\gamma)^{(D-1)/2}}\right\}.\end{aligned} \tag{C6}$$

For the coincidence case $\xi' = \xi$ from here one obtains

$$\begin{aligned}\mathcal{L}(\xi,\xi) = 2^{(D-5)/2}\pi m^{D+1}\xi^2\Gamma\left(\frac{D-1}{2}\right)\int_0^\infty \frac{dy}{\pi^2+y^2}\left\{[D+1-(D-1)\cosh y]\frac{K_{(D+1)/2}(z)}{z^{(D-1)/2}}\right. \\ \left.+(1-\cosh y)\frac{K_{(D-1)/2}(z)}{z^{(D-1)/2}}\right\}.\end{aligned} \tag{C7}$$

Comparing expressions (C4) and (C7) one obtains identity (53).

---

[1] V. M. Mostepanenko and N. N. Trunov, *The Casimir effect and its applications* (Oxford University Press, Oxford, 1997).
[2] G. Plunien, B. Muller, and W.Greiner, Phys. Rep. **134**, 87 (1986).
[3] K. A. Milton, in *Applied Field Theory*, Edited by C. Lee, H. Min, and Q-H. Park (Chungbum, Seul, 1999), p.1, hep-th/9901011; "Dimensional and Dynamical Aspects of the Casimir Effect: Understanding the Reality and Significance of Vacuum Energy", hep-th/0009173.
[4] N. D. Birrel and P. C. W. Davis, *Quantum fields in curved space* (Cambridge University Press, 1982).
[5] P. Candelas and D. Deutsch, Proc. Roy. Soc. Lond. A **354**, 79 (1977).
[6] J. Ambjørn and S. Wolfram, Ann. Phys. (N.Y.) **147** 1 (1983).
[7] C. M. Bender and K. A. Milton, Phys. Rev. D **50**, 6547 (1994); K. A. Milton, Phys. Rev. D **55**, 4940 (1997).
[8] A. Romeo and A. A. Saharian, "Casimir effect for scalar fields under Robin boundary conditions on plates," hep-th/0007242.
[9] A. A. Saharian, Phys. Rev. D **63**, 125007 (2001).
[10] A. Romeo and A. A. Saharian, Phys. Rev. D **63**, 105019 (2001).
[11] V. M. Mostepanenko and N. N. Trunov, Sov. J. Nucl. Phys. **42**, 812 (1985).
[12] S. N. Solodukhin, Phys. Rev. D **63** (2001) 044002.
[13] J. Ambjørn and S. Wolfram, Ann. Phys. (N.Y.) **147** 33 (1983).
[14] I.G. Moss, Class. Quantum Grav. **6**, 759 (1989).
[15] G. Esposito and A. Yu. Kamenshchik, Class. Quantum Grav. **12**, 2715 (1995).
[16] G. Esposito, A. Yu. Kamenshchik, and G. Polifrone, *Euclidean Quantum Gravity on Manifolds with Boundary* (Kluwer, Dordrecht, 1997).
[17] T. Ghergetta and A. Pomarol, Nucl. Phys. B **586**, 141 (2000).
[18] A. A. Saharian, Izv. AN Arm. SSR. Matematika **22**, 166 (1987) [Sov. J. Contemp. Math. Analysis **22**, 70 (1987)]; A. A. Saharian, Ph.D. thesis, Yerevan, 1987 (in Russian).
[19] A. A. Saharian, "The generalized Abel-Plana formula. Applications to Bessel functions and Casimir effect," Report No. IC/2000/14, hep-th/0002239.
[20] S. A. Fulling, Phys. Rev. D **7**, 2850 (1973).
[21] P. C. W. Davies, J. Phys. A **8**, 609 (1975).
[22] W. G. Unruh, Phys. Rev. D **14**, 870 (1976).
[23] P. Candelas and D. J. Raine, J. Math. Phys. **17**, 2101 (1976).

[24] *Integrals and series, Vol. 2: Special Functions,* edited by A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev (Harwood Academic, Newark, NJ, 1986).

[25] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical functions* (National Bureau of Standards, Washington D.C.,1964).

[26] A. Erdelyi, W. Magnus, F. Oberhettinger and F. G. Tricomi, *Higher Transcendental Functions* (McGraw-Hill, New-York, 1963).

[27] C. T. Hill, Nucl. Phys. B **277**, 547 (1986).

[28] K. Freese, C. T. Hill, and M. Mueller, Nucl. Phys. B **255**, 693 (1985).

[29] S. Tagaki, Prog. Theor. Phys. Suppl. **88**, 1 (1986).

[30] R. Jáuregui, M. Tarres, and S. Hacyan, Phys. Rev. D **43**, 3979 (1991).

[31] L. P. Chimento, A. E. Cossarini, and F. G. Pensa, Prog. Theor. Phys. **92**, 67 (1994).

[32] R. Medina and E. S. Moreira Jr, Phys. Rev. D **63**, 124022 (2001).

[33] H. H. Matevosyan and A. A. Saharian, "Casimir effect for massive scalar field under Robin boundary conditions," to be published.